



## King's Research Portal

DOI:

[10.1007/s00213-017-4542-8](https://doi.org/10.1007/s00213-017-4542-8)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Ettinger, U., Faiola, E., Kasparbauer, A. M., Petrovsky, N., Chan, R. C. K., Liepelt, R., & Kumari, V. (2017). Effects of nicotine on response inhibition and interference control. *Psychopharmacology*, 1093-1111. <https://doi.org/10.1007/s00213-017-4542-8>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**


Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**


If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**Figure 1: Effects on Saccade Task Latency and Error Rate**




*Note:* Figure (A) shows the main effect of Task Condition on saccadic latency, indicating shorter latencies for prosaccades than for antisaccades, and the interaction between Drug and Task Condition, indicating shorter mean latencies with nicotine compared to placebo for prosaccades (\* $p < .05$ ), but not for antisaccades ( $p > .05$ ). Figure (B) shows the main effect of Task Condition on error rate and the interaction between Order, Drug and Task Condition. Significantly fewer prosaccade errors occurred in the second compared to the first study session in the Nic-Plc group (\* $p < .05$ ). For further post-hoc tests, see main text. Error bars indicate the standard error.

**Figure 2: Effects on Stroop Task Reaction Time and Error Rate**




*Note:* Figure (A) shows the main effect of Task Condition, the interaction between Drug and Order, and the interaction between Drug, Task Condition and Order. In both groups, MRTs were shorter in the second study session compared to the first study session, but the size of this difference depended on Task Condition. Post-hoc t-tests indicated that in the Nic-Plc group, this difference was significant for incongruent (\* $p < .05$ ) and congruent (\* $p < .05$ ) but not for colour bar trials ( $p > .05$ ). For the Plc-Nic group, this difference was significant for incongruent trials (\* $p < .05$ ), but not for congruent and colour bar trials (both  $p > .05$ ). For further post-hoc tests, see main text. Error bars indicate the standard error. MRT is mean reaction time (in ms). Figure (B) shows the interaction effect between Drug and Order, indicating higher error rates in the second than the first study session. T-tests showed significant differences between error rates in the two drug conditions for the Plc-Nic group (\* $p < .05$ ) and, at trend level, for the Nic-Plc group ( $p = .05$ ). Error bars indicate the standard error. See main text for more details.

**Figure 3: Effects on Flanker Task Reaction Time**




*Note:* The figure shows the interaction between Drug and Order, indicating that MRTs were shorter in the second than the first study session, irrespective of Task Condition. These differences were significant both for the Nic-Plc group (\* $p < .05$ ) and for the Plc-Nic group (\* $p < .05$ ). The two Order groups did not differ in the placebo condition or nicotine conditions (both  $p > .05$ ). Error bars indicate the standard error. MRT is mean reaction time (in ms). See main text for more details.

**Figure 4: Effects on Shape Matching Task Reaction Time**




*Note:* The figure shows the interaction between Drug and Order, indicating that MRTs were shorter in the second than the first study session. This difference was significant for the Nic-Plc group ( $*p < .05$ ), but not the Plc-Nic group ( $p > .05$ ). The two order groups did not differ significantly in the placebo or nicotine conditions (both  $p > .05$ ). Error bars indicate the standard error. MRT is mean reaction time (in ms). See main text for more details.

**Figure 5: Effects on Simon Task Reaction Time**



*Note:* The figure shows the interaction between Drug and Task Condition. MRTs were higher with nicotine than placebo in incongruent (\* $p < .05$ ) but not congruent ( $p > .05$ ) trials, and in incongruent than congruent trials during both nicotine (\*\* $p < .001$ ) and placebo (\*\* $p < .001$ ). Error bars indicate the standard error. MRT is mean reaction time (in ms). See main text for more detail.

**Figure 6: Effects on Attention Network Test (ANT) Conflict Score**



*Note:* The figure shows the interaction between Drug and Order, indicating that conflict scores were lower in the second than the first study session. Differences were significant in each group (\* $p < .05$ ) and the two groups differed significantly under nicotine (\* $p < .05$ ) but not placebo ( $p > .05$ ). Error bars indicate the standard error. See main text for more detail.